arXiv:0908.0763v3 [quant-ph] 21 Dec 2009

Is the Quantum Zeno Effect Evolution's Choice for the Avian Compass?

I. K. Kominis
Department of Physics, University of Crete, Heraklion 71103, Greece and
Institute of Electronic Structure and Laser, Foundation for Research and Technology, Heraklion 71110, Greece

(Dated: December 21, 2009)

Magnetic-sensitive radical-ion-pair reactions are understood to underlie the biochemical magnetic compass used by avian species for navigation. Radical-ion-pair reactions were recently shown to manifest a host of quantum-information-science effects, like quantum jumps and the quantum Zeno effect. We here show that the quantum Zeno effect immunizes the magnetic and angular sensitivity of the avian compass mechanism against the deleterious and molecule-specific exchange and dipolar interactions.

PACS numbers: 82.20.-w

"In the history of natural selection, did nature ever come across a way to use quantum weirdness?" This is a question claimed [1] to have an affirmative answer, at least in regard with the apparent ability of photosynthetic antennae to efficiently guide the excitonic energy to the photosynthetic reaction center. The possibility of biological processes exhibiting non-trivial quantum effects, ordinarily thought to be suppressed in the decoherence-prone biological environment [2], has attracted an increasing attention in recent years. For example, significant experimental [3] and theoretical [4] progress has been recently made on elucidating the role of quantum coherence in the workings of photosynthetic antennae complexes. In a different front, radical-ion-pair reactions [5, 6] have been recently shown [7] to exhibit the full machinery of concepts and physical effects familiar from quantum information science. Radical-ion pairs play a fundamental role in a series of biologically relevant chemical reactions [8, 9], and are understood to underlie the biochemical magnetic compass used by avian species to navigate in earth's magnetic field [10], as corroborated by several recent experiments [11].

Radical-ion pairs are formed by a charge transfer following a photoexcitation of a donor-acceptor dyad, leading to two molecular ions and two unpaired electrons. The latter can either be in the spin singlet or in the spin triplet state. Magnetic interactions with the external magnetic field and hyperfine interactions with the molecule's magnetic nuclei bring about a coherent singlet-triplet oscillation. At some random instant in time the reaction is terminated, since the radical-ion-pair undergoes charge recombination, leading to the reaction products. Spin conservation enforces spin selectivity of the recombination process, i.e. singlet (triplet) radical-ion pairs recombine to singlet (triplet) neutral products. Moreover, anisotropic hyperfine interactions within the molecule render the reaction yields dependent on the inclination of the external magnetic field with respect to a molecule-fixed coordinate frame. The avian magnetoreception mechanism exactly rests on both aforementioned aspects, i.e. the reaction yield's magnetic and angular sensitivity.

Interestingly, intra-molecule magnetic interactions are more complicated. Both spin-exchange and long-range dipolar interactions affect reaction dynamics, to an extent dependent on the particular molecular structure. In this respect, it has been recently shown [12] that the presence of exchange and/or dipolar interactions significantly suppresses the magnetic and angular sensitivity of the reaction yields, thus severely degrading the mechanism's functionality. Along the same lines, it was concluded [12] that only when the molecular parameters determining J and D, the exchange and dipolar couplings, are fine-tuned so that the effects of these two interactions cancel each other, is the magnetic and angular sensitivity of the reaction restored. Although a remarkable result, it is hard to imagine that Nature has conjured up such a fortuitous cancellation for a functionally important biological sensor. Even more so since J and D depend exponentially on molecule-specific parameters, like the donor-acceptor distance.

However, it was recently shown [7] that the fundamental description of radical-ion-pair reactions following from quantum measurement theory reveals the actual presence of non-trivial quantum effects, such as the quantum Zeno effect [13]. We will here show that when the quantum Zeno effect is manifested, the reaction's magnetic and angular sensitivity is practically independent of the presence or not of exchange and/or dipolar interactions. This realization has profound implications for the robustness of this biological sensor, i.e. a non-trivial quantum effect renders the sensor insensitive to molecule-specific parameters [12], such as donor-acceptor distance (affecting the exchange coupling and the long-range dipolar coupling) and the inter-radical medium and the particular electronic structure (affecting the exchange coupling).

The spin state of the radical-ion-pair is described by a $4n$-dimensional density matrix ρ , where the factors 4 and n are the spin multiplicities of the two electrons and the molecule's nuclear spins, respectively. The unconditional evolution of ρ is given [7] by the Liouville equation

$$d\rho/dt = -i[\mathcal{H}_m, \rho] - (k_S + k_T)(Q_S\rho + \rho Q_S - 2Q_S\rho Q_S) \tag{1}$$

where Q_S and Q_S are the singlet and triplet state projection operators, respectively. The first term in (1) is the unitary evolution due to the magnetic interactions, whereas the second term describes the measurement-induced evolution due to the singlet (rate k_S) and triplet (rate k_S) recombination channel. The reaction is terminated at a random instant in time according to the quantum jump equations [14]

$$\begin{aligned} dp_S &= 2k_S\langle Q_S\rangle dt \\ dp_T &= 2k_T\langle Q_T\rangle dt \end{aligned} \tag{2}$$

which give the probability of singlet and triplet recombination taking place within the time interval between t and $t + dt$.

The magnetic Hamiltonian $\mathcal{H}_m = \mathcal{H}_Z + \mathcal{H}_{hf} + \mathcal{H}_{ex}$ is composed of $\mathcal{H}_Z$, the Zeeman interaction of the two unpaired electrons (nuclear Zeeman interaction is negligible) with the external magnetic field, $\mathcal{H}_{hf}$, the hyperfine couplings of the electrons with the surrounding nuclear spins, and finally the spin-exchange interaction, $\mathcal{H}_{ex}$. For the transparency of the following discussion we will ignore the dipolar interaction as its inclusion leads to exactly the same conclusions. The Zeeman interaction Hamiltonian that will be used for the study of the magnetic sensitivity is $\mathcal{H}_{Z,magn} = \omega(s_{1z} + s_{2z})$, where the magnetic field of magnitude B is assumed to be in the z-axis ($\omega = \gamma B$, with γ=1.4 MHz/G). For the study of the angular sensitivity we take the magnetic field, again of magnitude B, to be in the x-y plane, hence the Zeeman interaction term will be $\mathcal{H}_{Z,ang} = \omega\cos\phi(s_{1x} + s_{2x}) + \omega\sin\phi(s_{1y} + s_{2y})$. In the following we will consider the simplest physically realizable radical-ion-pair containing just one spin-1/2 nucleus (in which case dim(ρ)=8), hence the hyperfine interaction Hamiltonian is $\mathcal{H}_{hf} = \mathbf{I}\cdot\mathbf{A}\cdot\mathbf{s}_1$, where $\mathbf{A}$ is the hyperfine coupling tensor of the single nuclear spin $\mathbf{I}$ existing in e.g. the donor molecule with the donor's unpaired electron. We will consider the simplest case where the hyperfine tensor is diagonal with one non-zero component $A_{xx} = a$, to provide for the angular sensitivity on the $x-y$ plane. Thus $\mathcal{H}_{hf} = as_{1x}I_x$. Finally the spin-exchange Hamiltonian is $\mathcal{H}_{ex} = J\mathbf{s}_1\cdot\mathbf{s}_2$. In the following we will calculate the magnetic and angular sensitivity of the reaction for two regimes: (i) the "traditional" regime where both recombination rates k_S and k_T are equal and small compared to the magnetic frequency scale h (set by a and ω), i.e. $k_S = k_T \leq a, \omega$. It is in this regime that almost all calculations have been performed based on the previous, phenomenological theory [7]. We then study the regime (ii) where $k_T \gg k_S$ and $k_T \sim \omega$, i.e. when the quantum Zeno effect is manifested, since the total measurement rate $k = k_S + k_T$ is dominated by the triplet recombination channel, strongly re-projecting the radical-ion-pair's spin state to the singlet state, from which it is considered to start out, i.e. the initial condition is $\rho(t = 0) = Q_S/2$ (which satisfies $\mathrm{Tr}\{\rho\} = 1$ and $\mathrm{Tr}\{\rho \mathrm{Q_S}\} = 1$). We will show that in the regime (i) the inclusion of the spin exchange interaction indeed degrades the magnetic and angular sensitivity of the reaction, as has already been analyzed [12]. However, regime (ii) exhibits a significant magnetic and angular sensitivity with their dependence on the exchange coupling J being almost entirely suppressed.

In Fig. 1 (insets) we show the triplet yield Y_T of the reaction as a function of the external magnetic field (using $\mathcal{H}_{Z,magn}$) and the field's angle (using $\mathcal{H}_{Z,ang}$). The triplet yield is calculated from

$$Y_T[\%] = \frac{100}{N_0}\int_0^\infty dp_T(t)N(t) \tag{3}$$

where N_0 is the initial number of radical-ion pairs (starting out in the singlet state) and $N(t)$ the remaining number of radical-ion pairs at time t, evolving as [7] $dN = -N(dp_S + dp_T)$. That is, in the time interval between t and $t+dt$ the number of created triplet products will be the number of radical-ion-pairs existing up to that time, $N(t)$, times the probability dp_T for triplet-channel recombination. This is summed in (3) until all molecules have recombined one way or another. The probabilities dp_S and dp_T are calculated from (2), whereas the expectation values $\langle Q_S\rangle$ and $\langle Q_T\rangle$ entering (2) are obtained by evolving the density matrix according to (1). The magnetic sensitivity of the reaction at earth's field of interest for the avian compass is just the slope dY_T/dB calculated at B=0.5 G. The smallest measurable change of the magnetic field (magnetic precision), δB, and the precision $\delta\phi$ of the determination of the magnetic field angle with respect to the molecule's x-axis (angular precision or heading error) both follow from the previous calculations if the smallest measurable reaction yield change, δY_T, is known. It thus follows that

$$\delta B = \frac{\delta Y_T}{|dY_T/dB|_{B=0.5\ \mathrm{G}}} \tag{4}$$

$$\delta\phi = \frac{\delta Y_T}{(Y_{T,max} - Y_{T,min})/\Delta\phi} \tag{5}$$

where $Y_{T,min}$ and $Y_{T,max}$ are the minimum and maximum values of the yield $Y_T(\phi)$ and $\Delta\phi = 90^{\mathrm{o}}$ is the angular width of the full swing between $Y_{T,min}$ and $Y_{T,max}$. To make further progress towards the absolute magnetic and angular sensitivity of the avian compass, the smallest "measurable" change in the reaction yield δY_T must be known or estimated. Obviously δY_T depends on the particular realization of the biochemical mechanism transducing the radical-ion-pair reaction yield to a physiological signal. On rather general grounds it has been shown [15] that δY_T is connected to N_R, the number of neuronal receptors sensitive to the radical-ion-pair



FIG. 1: Magnetic and angular precision. For all calculations the hyperfine coupling is a =5 G. (a) Inset shows an example of the magnetic sensitivity of the triplet yield plotted in the quantum Zeno regime for two values of the exchange coupling J. For higher (lower) values of the exchange coupling the plots are similar but appear at a lower (higher) offset. The yellow band is the 0.05% error band corresponding to the measurement precision of the reaction yield. Main graph shows the magnetic precision, δB, extracted from the slope at B =0.5 G of plots like the ones in the inset, as a function of the exchange coupling J for the two regimes of the recombination rates. (b) Similar calculation as in (a), but using $\mathcal{H}_{Z,ang}$ instead of $\mathcal{H}_{Z,magn}$, in order to derive the angular precision or heading error.

reaction product molecules, by $(\delta Y_T)^2 = 4/N_R$. We choose $N_R = 1.6 \times 10^7$, in order to set δY_T at the value $\delta Y_T = 0.05\%$. The chosen value of N_R and hence δY_T is realistic [15] and has the consequence that it sets the magnetic precision at zero exchange coupling (shown in Fig. 1a) at the value of $\delta B \approx 0.01$ G, i.e. at 2% of earth's field. This level of magnetic precision is understood [10] to be actually the case for several avian species. It is stressed, however, that the following considerations are qualitatively independent of the particular value of δY_T, which just sets the absolute scale of the derived magnetic and angular precision. It is clearly seen in Fig. 1a that in the traditional regime (i) the magnetic precision plunges from $\delta B \approx 0.01$ G at $J = 0$ to over 1 G at J=15 n G. Even at J=8 G the change of earth's field is barely measurable. On the contrary, in the quantum Zeno regime (ii) the magnetic precision drops only to $\delta B \approx 0.07$ G at J=15 G. Similarly, as shown in Fig. 1b, the angular precision in regime (i) drops dramatically with increasing J, with a complete loss of heading information for J=10 G or higher. In contrast, in the quantum Zeno regime the angular precision of about $\delta\phi = 20^o$ at the highest value of the exchange coupling is actually at the level of experimental observations [16] of the heading error of the avian compass. At the same time the heading error is practically unaffected by the presence or not of exchange and dipolar interactions, varying between $\delta\phi = 3.5^o$ and $\delta\phi = 20.5^o$ in the full span of the exchange coupling J.

We will now explain the robust magnetic and angular precision in the quantum Zeno regime. This follows by considering the behavior of the eigenvalues of the master equation (1), which are obtained by diagonalizing the matrix $\mathcal{M}$ (of dimension $\dim(\rho)^2$) that satisfies $d\tilde{\rho}/dt = \mathcal{M}\tilde{\rho}$, where $\tilde{\rho}$ is a column matrix containing all matrix elements of ρ . The resulting eigenvalues are of the form $-\lambda + i\Omega$, with $\lambda \geq 0$ being the decay rate and Ω the oscillation frequency of the particular eigenmode. As is in general the case with the quantum Zeno effect [17], some of the eigenvalues have decay rates increasing with the measurement rate k as $\lambda \sim k$, while the others (responsible for the quantum Zeno effect) decrease with k as $\lambda_{qZ} \sim h^2/k$, where h is the characteristic frequency scale of the system, here determined by the magnetic Hamiltonian $\mathcal{H}_m$.

Now, the exchange Hamiltonian can be written (up to an additive constant) as $\mathcal{H}_{ex} = -JQ_S$. Furthermore, as is known from quantum measurement theory [18], the deterministic evolution of the system's quantum state (due to the unitary Hamiltonian evolution and the measurement of Q_S with rate $k = k_S + k_T$) is generated by the non-hermitian operator $\mathcal{K} = \mathcal{H}_m - ikQ_S$. It is easily seen that if $\mathcal{H}_m^{J=0}$ is the magnetic Hamiltonian without the exchange interaction $\mathcal{H}_{ex}$, then $\mathcal{K} = \mathcal{H}_m^{J=0} - i(k - iJ)Q_S$, i.e. the inclusion of the exchange interaction is equivalent to replacing k with with an imaginary measurement rate $k - iJ$. We can now complete the argument as follows: the eigenvalues with a real part that scales as $\lambda \sim k$ pick up an oscillation frequency (in addition to Ω) of $-J$, the effect of which roughly averages out. On the other hand, the eigenvalues with the quantum Zeno scaling $\lambda_{qZ} \sim h^2/k$ suffer a change in their real part which becomes (since in our case $k/J \ll 1$) $\lambda'_{qZ} \approx h^2k/J^2 \ll \lambda_{qZ}$. Thus the spin state evolution is slowed down, the radical-ion-pair is projected stronger to the singlet state leading to an increased singlet yield, or equivalently a decreased triplet yield, which is exactly what is observed in Fig. 1 (insets). The above argument is pictured in Fig. 2a, which shows the time evolution of $N(t)$, the number of existing (non-reacted) radical-ion pairs. The reaction is considered to be terminated when $N(t)/N(0) \approx 5 \times 10^{-4}$, i.e. when the reaction yield is known to within δY_T. It is clearly seen that in the quantum Zeno regime, the reaction time de-



FIG. 2: Reaction Dynamics. (a) Time evolution of the number of radical ion pairs for the two regimes of the recombination rates, for different values of J. The dashed line at $N = 5 \times 10^6$ signifies the "termination" of the reaction, i.e. the point when the remaining radical-ion pairs are 0.05% of the initial number $N(t = 0) = 10^{10}$. (b) Time evolution of $\langle Q_S(t)\rangle$ and $\langle Q_T(t)\rangle$ for B=0.5 G and J=10 G. (c) Triplet yield as a function of the magnetic field for J=10 G.

pends on J in the way outlined before. On the contrary, when $k_S = k_T = \kappa$, the change in the number of radical-ion-pairs during the time interval dt is ($Q_S + Q_T = 1$) $dN = -Ndt(2k_S\langle Q_S\rangle + 2k_T\langle Q_T\rangle) = -2\kappa dtN$, hence $N(t) = N(0)e^{-2\kappa t}$, i.e. the reaction time is proportional to $1/\kappa$ and independent of J. The result is that during the short reaction time in the traditional regime, the triplet probability Q_T has not increased appreciably (Fig. 2b), and the triplet yield is small, as shown in Fig. 2c. In contrast, in the quantum Zeno regime the reaction has enough time to "sample" large values of Q_T and lead to a triplet yield about an order of magnitude higher, hence the higher sensitivity in this regime. In other words, as seen in Fig. 2c, the relative change $\delta Y_T/Y_T$ of the triplet yield with the magnetic field is roughly the same in both cases, but the absolute value of Y_T differs by a factor of 10, leading to respectively high slopes dY_T/dB and $dY_T/d\phi$. To summarize, the quantum measurement dynamics inherent in the recombination process of radical-ion pairs result in "delocalization" of the electron spin state at long times, as evidenced in Fig. 2b. The asymmetric ($k_T \gg k_S$) recombination rates result in the actual manifestation of the quantum Zeno effect and the J-dependence of the reaction dynamics. The interplay of these two effects provides for the robust magnetic and angular sensitivity in the presence of the exchange interaction.

Whether Nature has engineered molecules realizing a fine-tuned cancellation [12] of the adverse effects of exchange and dipolar interactions, or on the other hand, has evolved the avian compass into operating at the quantum Zeno regime, remains to be discovered.

[1] S. Lloyd, Nature Phys. **5**, 164 (2009).
[2] P. C. W. Davies, Biosystems **78**, 69 (2004); D. Abbott *et al.*, Fluctuations and Noise Letters **8**, C5 (2008).
[3] G. S. Engel *et al.*, Nature **446**, 782 (2007); H. Lee, Y. C. Cheng and G. R. Fleming, Science **316**, 1462 (2007).
[4] M.B. Plenio and S.F. Huelga, New. J. Phys. **10**, 113019 (2008); M. Mohseni, P. Rebentrost, S. Lloyd and A. Aspuru-Guzik, J. Chem. Phys. **129**, 174106 (2008).
[5] K. Schulten, Adv. Solid State Phys. **22**, 61 (1982).
[6] U. Steiner and T. Ulrich, Chem. Rev. **89**, 51 (1989).
[7] I. K. Kominis, Phys. Rev. E. **80**, 056115 (2009).
[8] S. G. Boxer, C. E. D. Chidsey and M. G. Roelofs, Ann. Rev. Phys. Chem. **34**, 389 (1983).
[9] C. R. Timmel, U. Till, B. Brocklehurst, K. A. McLauchlan and P. J. Hore, Molec. Phys. **95**, 71 (1998); C. R. Timmel, F. Cintolesi, B. Brocklehurst and P. J. Hore, Chem. Phys. Lett. **334**, 387 (2001); C. R. Timmel and K. B. Henbest, Phil. Trans. R. Soc. Lond. A **362**, 2573 (2004).
[10] T. Ritz, S. Adem and K. Schulten, Biophys. J. **78**, 707 (2000); W. Wiltschko and R. Wiltschko, J. Comp. Physiol. A **191**, 675 (2005).
[11] T. Ritz, P. Thalau, J. B. Phillips, R. Wiltschko and W. Wiltschko, Nature **429**, 177 (2004); K. Maeda *et al.*, Nature **453**, 387 (2008); T. Ritz *et al.*, Biophys. J. **96**, 3451 (2009); Zapka, M. *et al.*, Nature **461**, 1274 (2009).
[12] O. Efimova and P. J. Hore, Biophys. J. **94**, 1565 (2008).
[13] P. Facchi and S. Pascazio, Fortschr. Phys. **49**, 941 (2001); K. Koshino and A. Shimizu, Phys. Rep. **412**, 191 (2005); A. G. Kofman and G. Kurizki, Nature **405**, 546 (2000).
[14] H. M. Wiseman, Quantum Semiclass. Opt. **8**, 205 (1996).
[15] J. C. Weaver, T. E. Vaughan and E. R. Astumian, Nature **405**, 707 (2000).
[16] W. Wiltschko, R. Wiltschko, J. Exp. Biol. **199**, 29 (1996); W. W. Cochran, H. Mouritsen and M. Wikelski, Science **304**, 405 (2004).
[17] E. W. Streed *et al.*, Phys. Rev. Lett. **97**, 260402 (2006).
[18] H. -P. Breuer and F. Petruccione, *The theory of open quantum systems* (Oxford University Press, 2002).